

IMPORTANT INFORMATION FOR AFTER YOUR SURGERY

Pickerington Manor Jaw & Facial Surgery



ORAL CARE & POSTOPERATIVE INSTRUCTIONS

After Implants/Bone Grafting

DR. M. BATH

*Practice Limited to
Oral and Maxillofacial Surgery
Dental Implant Surgery*

**1575 CROSS CREEKS BLVD.
PICKERINGTON, OHIO 43147**

614-751-7500

www.pickeringtonoms.com

**IF AT ANY TIME YOU ARE UNCERTAIN ABOUT YOUR TREATMENT
OR THESE INSTRUCTIONS PLEASE CALL.**

IMPORTANT INFORMATION FOR AFTER YOUR SURGERY

GENERAL INFO

YOU HAVE JUST UNDERGONE A SURGICAL PROCEDURE. VARYING DEGREES OF PAIN MAY CONTINUE FOR ABOUT ONE WEEK FOLLOWING SURGERY. YOU MAY ALSO EXPERIENCE DISCOMFORT IN THE EARS, THE THROAT, THE SIDE OF THE HEAD, AND UNDER YOUR LOWER JAW. THERE WILL BE AN OPENING WHERE THE TOOTH WAS REMOVED. PLEASE TRY TO KEEP YOUR FINGERS AND TONGUE OUT OF THIS AREA. TRY TO **LIMIT YOUR ACTIVITIES** AS MUCH AS POSSIBLE ON THE DAY OF SURGERY TO REDUCE BLEEDING AND PERMIT FORMATION OF A CLOT IN THE WOUND. THIS IS NECESSARY FOR PROPER HEALING. SURGICAL WOUNDS OF THE MOUTH USUALLY HEAL QUICKLY AND WITHOUT COMPLICATIONS. IF YOU TAKE THE SIMPLE PRECAUTIONS DESCRIBED IN THIS PAMPHLET, IT WILL LIKELY MAKE YOUR RECOVERY AN EASIER ONE.

SWELLING

SOME SWELLING IS TO BE EXPECTED, AND SHOULD NOT ALARM YOU. THE MOST SWELLING USUALLY OCCURS DURING THE FIRST 48 HOURS, AND THEN BEGINS TO DECREASE. APPLYING ICE TO THE AREA FOR THE FIRST 24 HOURS, REPEATEDLY, WILL HELP REDUCE THE AMOUNT OF SWELLING. KEEPING YOUR HEAD ELEVATED DURING THIS PERIOD CAN ALSO HELP LIMIT YOUR SWELLING. THE AMOUNT OF SWELLING VARIES FROM PERSON TO PERSON AND DEPENDS UPON THE COMPLEXITY OF YOUR SURGERY. IF YOUR SWELLING PERSISTS OR INCREASES, AFTER THE FIRST 3 TO 4 DAYS, CALL OUR OFFICE. IF A VEIN USED TO ADMINISTER GENERAL ANESTHESIA BECOMES SORE AND SWOLLEN, PLEASE NOTIFY OUR OFFICE IMMEDIATELY.

PAIN AND MEDICATIONS

WE WISH TO INFORM YOU OF OUR OFFICE POLICY: NO PAIN MEDICATIONS

CAN BE CALLED IN AFTER OFFICE HOURS. MOST OFTEN, YOU WILL EXPERIENCE SOME DISCOMFORT FOR A FEW DAYS FOLLOWING YOUR PROCEDURE. THE **PRESCRIPTION FOR PAIN SHOULD BE FILLED ONLY IF NEEDED**, AND TAKEN AS DIRECTED. IF YOU WERE NOT PRESCRIBED ANY MEDICATION OR YOU HAVE COMPLETED YOUR PRESCRIPTION, TAKE OVER THE COUNTER MEDICATIONS SUCH AS **ADVIL** (IBUPROFEN), **ALEVE** (NAPROSYN), OR **TYLENOL** (ACETOMENOPHEN) AS DIRECTED ON THE BOTTLE. **DO NOT TAKE ANY OTHER PAIN MEDICINES OR SEDATIVES WITH YOUR PRESCRIBED DRUGS,** UNLESS YOUR DOCTOR HAS SPECIFICALLY DIRECTED YOU TO DO SO.

NARCOTICS: IT CAN BE EXTREMELY DANGEROUS TO CONSUME ALCOHOL WHILE YOU ARE TAKING NARCOTIC PAIN MEDICATIONS. ALSO, SINCE THESE DRUGS CAN CAUSE DROWSINESS OR ALTER YOUR PERCEPTION IT IS ADVISED YOU **SHOULD NOT TAKE THEM IF YOU ARE DRIVING ANY MOTOR VEHICLE, OPERATING MACHINERY, OR DOING ANYTHING THAT REQUIRES ALERTNESS.**

INCREASING PAIN ON THE THIRD TO FIFTH DAY AFTER TOOTH REMOVAL MAY INDICATE A PROBLEM WITH HEALING. THIS PAIN IS USUALLY SEVERE AND UNRESPONSIVE TO PAIN MEDICATIONS. IF THIS HAS OCCURRED, CONTACT THE OFFICE IMMEDIATELY FOR AN APPOINTMENT.

ANTIBIOTICS: WHEN ANTIBIOTICS ARE PRESCRIBED, PLEASE TAKE THEM AS DIRECTED AND COMPLETE THE ENTIRE AMOUNT THAT WAS GIVEN. IF YOU DEVELOP A RASH, HIVES OR PERSISTANT ITCHING, YOU MAY BE HAVING AN ALLERGIC REACTION. STOP TAKING THE MEDICATION AND CALL THE OFFICE. IF YOU **EXPERIENCE TIGHTNESS IN YOUR CHEST OR DIFFICULTY BREATHING, SEEK EMERGENCY MEDICAL CARE.**

BIRTH CONTROL USERS: PLEASE USE ALTERNATIVE MEANS OF CONTRACEPTION AND TAKE EXTRA PRECAUTIONS. THE USE OF PRESCRIPTION MEDICATIONS AND ANTIBIOTICS CAN CAUSE YOUR BIRTH CONTROL **TO BE LESS EFFECTIVE.**

IMPORTANT INFORMATION FOR AFTER YOUR SURGERY

BLEEDING

SOME OOZING FROM THE SURGICAL SITE MAY CONTINUE FOR THE FIRST 24 HOURS. HOWEVER, IF THERE IS MORE THAN SLIGHT OOZING, THE FOLLOWING PROCEDURES CAN HELP **CONTROL THE BLEEDING**:

1. USE CLEAN GAUZE TO **GENTLY** REMOVE THE BLOOD FROM YOUR MOUTH AND AROUND SURGICAL AREAS. ALL BLOOD CLOTS THAT ARE **NOT IN THE WOUND** SHOULD BE REMOVED. DO NOT ATTEMPT TO REMOVE ANYTHING FROM INSIDE THE SOCKET OR INCISION AREA.
2. PLACE A CLEAN FOLDED GAUZE PAD ON THE BLEEDING AREA / SOCKET. NOT JUST BETWEEN YOUR TEETH.
3. CLOSE YOUR MOUTH TIGHTLY OVER THE GAUZE PAD SO THAT THERE IS **PRESSURE AGAINST THE SPOT THAT IS BLEEDING**. MAKE SURE THE GAUZE DOES NOT END UP BETWEEN YOUR TEETH, AS IT IS CRUCIAL TO PUT PRESSURE DIRECTLY OVER THE SITE THAT IS BLEEDING.
4. MAINTAIN THE PRESSURE FOR ABOUT **20 TO 30 MINUTES**. RESIST THE URGE TO CHECK ON THE SITE PRIOR TO THIS PERIOD OF TIME. REPEAT THE PROCESS IF NECESSARY.
5. **AVOID SPITTING** AND RINSING IF BLEEDING SUCH AS THIS OCCURS. DOING SO WILL CAUSE THE BLEEDING TO START AGAIN, OR CONTINUE.
6. IF THE BLEEDING HAS NOT COME UNDER CONTROL AFTER A COUPLE OF ATTEMPTS, A **MOISTENED TEA BAG** CAN BE PLACED OVER THE SITE TO AIDE IN THE CLOTTING PROCESS OR YOU MAY CALL DR. BATH.

SUTURES

SUTURES (STITCHES) MAY HAVE BEEN USED DURING YOUR SURGERY. THEY WILL **USUALLY DISSOLVE OR FALL OUT WITHIN A WEEK**. IF THEY ARE BOTHERSOME, THEY CAN BE REMOVED AT YOUR FOLLOW UP APPOINTMENT. OCCASSIONALLY LONGER LASTING SUTURES ARE USED. YOU WILL BE INFORMED IF THIS APPLIES TO YOU. PLEASE DO NOT PICK OR PULL AT YOUR STITCHES. TRY TO LEAVE THE AREA ALONE, AS **A PROBING TONGUE, AND PULLING OF CHEEKS OR LIPS CAN DO MUCH DAMAGE TO SURGICAL AREAS**.

ORAL HYGIENE

AFTER SURGERY, **WAIT UNTIL THE FOLLOWING DAY** TO START YOUR ORAL CARE. BEGIN RINSING YOUR MOUTH WITH WARM SALT WATER (ONE-HALF TEASPOON OF SALT IN AN 8 OZ. GLASS OF WARM WATER) AT LEAST FIVE TO SIX TIMES A DAY. IT IS IMPORTANT TO CONTINUE WITH THE EVERYDAY BRUSHING AND HYGIENE OF YOUR MOUTH. WHEN BRUSHING, BE GENTLE NEAR THE SURGICAL SITES AND STITCHES. YOU ARE LESS LIKELY TO GET A POSTOPERATIVE INFECTION IF YOUR MOUTH AND TEETH ARE KEPT CLEAN. AFTER YOUR FOLLOW-UP APPOINTMENT, YOU WILL LIKELY BE STARTED ON A MEDICATED RINSE. PLEASE WAIT TO USE ANY ALCOHOL-BASED RINSES.

DIET

DURING THE FIRST FEW DAYS: AVOID HOT TEMPERATURE FOODS FOR THE FIRST 24 HOURS, AVOID DRINKING THROUGH A STRAW, AVOID SPICY FOODS AND CARBONATED BEVERAGES. NOT DOING SO COULD AFFECT YOUR HEALING OR CAUSE MORE BLEEDING. AFTER A **LOCAL ANESTHETIC**, YOU SHOULD TAKE CARE UNTIL THE NUMBNESS HAS WORN OFF. AFTER A

IMPORTANT INFORMATION FOR AFTER YOUR SURGERY

GENERAL ANESTHETIC, DO THE SAME AS WITH A LOCAL, AND START OUT WITH CLEAR FLUIDS AT FIRST AND GRADUALLY WORK UP TO A SOFT DIET. **AFTER SURGERY STAY ON A SOFT NO-CHEW DIET**, UNTIL YOU ARE INSTRUCTED OTHERWISE (Can be 2-4 months or longer). **TONGUE PROBING OR CHEWING ON AN IMPLANT OR BONEGRAFT CAN CAUSE THE SURGERY TO FAIL AND REQUIRE FURTHER TREATMENT.** EAT NUTRITIOUS FOODS AND EAT SMALL AMOUNTS AS FREQUENTLY AS YOU CAN DURING THE FIRST FEW DAYS. IF YOU FIND THAT YOU CANNOT KEEP YOUR FOOD DOWN, PLEASE CALL THE OFFICE FOR ADVICE, OR AN ANTI-NAUSEA PRESCRIPTION IF NEED BE.

IMPLANTS & BONE GRAFTS

ORAL CARE, IF YOU HAVE HAD IMPLANTS OR BONE GRAFTS, WILL BE SIMILAR TO OTHER ORAL SURGERY PATIENTS. IN ADDITION, YOU MAY OBSERVE SOME OF THE FOLLOWING CONDITIONS: SMALL (SAND-LIKE) BONE PARTICLES IN THE GRAFT SIGHT. INCREASED SWELLING & TIGHTNESS IN THE MUSCLES OF YOUR JAW MAY CAUSE DISCOMFORT OR DIFFICULTY WITH OPENING YOUR MOUTH. YOUR BITE MAY FEEL "OFF," ADJACENT TEETH MAY ACHE, FEEL LOOSE OR BE TEMPERATURE SENSITIVE. **PROLONGED NUMBNESS OR TINGLING IN YOUR LIP OR TONGUE**, THOUGH RARELY PERMANENT, MAY LAST SEVERAL WEEKS OR MONTHS. A **SLIGHT FEVER** FOR THE FIRST 48 HOURS IS NOT UNCOMMON, BUT IF IT PERSISTS, PLEASE CALL.

SMOKING

SMOKING IS NOT ADVISED AFTER IMPLANT OR GRAFTING SURGERY. SMOKING DELAYS HEALING AND CAN RESULT IN INCREASED PAIN, A DRY SOCKET, OR **FAILURE OF YOUR IMPLANT OR BONE GRAFT TO HEAL.** NOT SMOKING IS A DIFFICULT RULE TO FOLLOW, BUT EVEN THE SLIGHTEST BIT CAN COMPLICATE YOUR RECOVERY. **THE INVESTMENT YOU HAVE MADE IN RESTORING YOUR MOUTH AND JAWS COULD BE WASTED.** IF YOU NEED ASSISTANCE IN SMOKING CESSATION, PLEASE ASK.

UNUSUAL CONDITIONS

PLEASE CALL THE OFFICE IF ANY OF THE FOLLOWING CONDITIONS ARE NOTED:

1. FEVER OVER 101 DEGREES.
2. SEVERE, BRIGHT RED BLEEDING THAT HAS NOT RESPONDED TO THE INSTRUCTIONS IN THIS PAMPHLET.
3. SEVERE OR LARGE HARD SWELLING AFTER THE 4TH POSTOPERATIVE DAY.
4. RASH, DIFFICULTY BREATHING, SEVERE VOMITING, OR ANY OTHER SUSPECTED REACTION TO YOUR MEDICATION(S).

PLEASE BE ADVISED

THE DOCTOR CAN BE REACHED AFTER HOURS FOR EMERGENCIES, BY CALLING THE OFFICE NUMBER ON THE FRONT, AND A CONTACT NUMBER WILL BE GIVEN DURING AFTERHOUR CALLS. IF YOU ARE HAVING DIFFICULTIES DURING THE DAY, WHEN AT ALL POSSIBLE, WE REQUEST YOU CALL DURING OFFICE HOURS TO ENSURE PROMPT ASSISTANCE WITH YOUR CONCERNS.